

Exercice 1 *Intégration directe*

1. $\int \sqrt{2px} dx = \frac{2x}{3} \sqrt{2px}$
2. $\int x(x+a)(x+b) dx = \frac{x^4}{4} + \frac{(a+b)x^3}{3} + \frac{abx^2}{2}$
3. $\int \left(a^{\frac{2}{3}} - x^{\frac{2}{3}}\right)^3 dx = a^2x - \frac{9}{5}a^{4/3}x^{5/3} + \frac{9}{7}a^{2/3}x^{7/3} - \frac{x^3}{3}$
4. $\int \tan^2(x) dx = \tan(x) - x$
5. $\int \frac{dx}{x^2 - 10} = \frac{1}{2\sqrt{10}} \ln \left| \frac{x - \sqrt{10}}{x + \sqrt{10}} \right|$
6. $\int 3^x e^x dx = \frac{3^x e^x}{1 + \ln(3)}$.

Exercice 2 *Changement de variable*

1. $\int \frac{\sqrt{x} + \ln(x)}{x} dx = 2\sqrt{x} + \frac{\ln^2(x)}{2}$
2. $\int \frac{dx}{(a+b) - (a-b)x^2} = \frac{1}{2\sqrt{a^2 - b^2}} \ln \left| \frac{\sqrt{a+b} + x\sqrt{a-b}}{\sqrt{a+b} - x\sqrt{a-b}} \right|$ pour $0 < b < a$
3. $\int \frac{x^2 + 5x + 7}{x + 3} dx = \frac{x^2}{2} + 2x + \ln|x + 3|$
4. $\int \frac{3x + 1}{\sqrt{5x^2 + 1}} dx = \frac{3}{5} \sqrt{5x^2 + 1} + \frac{1}{\sqrt{5}} \ln(x\sqrt{5} + \sqrt{5x^2 + 1})$
5. $\int \frac{xdx}{\sqrt{a^4 - x^4}} = \frac{1}{2} \arcsin\left(\frac{x^2}{a^2}\right)$
6. $\int \sqrt{\frac{\arcsin(x)}{1 - x^2}} dx = \frac{2}{3} \sqrt{\arcsin^3(x)}$
7. $\int \frac{dx}{\sqrt{(1+x^2) \ln(x + \sqrt{1+x^2})}} = 2\sqrt{\ln(x + \sqrt{1+x^2})}$
8. $\int \sqrt{1 + 3 \cos^2(x)} \sin(2x) dx = -\frac{2}{9} \sqrt{(1 + 3 \cos^2(x))^3}$
9. $\int \frac{x^3 dx}{x^8 + 5} = \frac{1}{4\sqrt{5}} \arctan\left(\frac{x^4}{\sqrt{5}}\right)$
10. $\int \frac{e^{\arctan(x)} + x \ln(1+x^2) + 1}{1+x^2} dx = e^{\arctan(x)} + \frac{\ln^2(1+x^2)}{4} + \arctan(x)$
11. $\int \frac{\sin(x) - \cos(x)}{\sin(x) + \cos(x)} dx = -\ln|\sin(x) + \cos(x)|$
12. $\int \frac{\arcsin(x) + x}{\sqrt{1-x^2}} dx = \frac{\arcsin^2(x)}{2} - \sqrt{1-x^2}$

Exercice 3 *Intégration par parties*

1. $\int \frac{\ln(x)}{\sqrt{x}} dx = 2\sqrt{x} \ln(x) - 4\sqrt{x}$
2. $\int (x^2 + 5x + 6) \cos(2x) dx = \frac{2x^2 + 10x + 11}{4} \sin(2x) + \frac{2x + 5}{4} \cos(2x)$
3. $\int x^2 \ln(x) dx = \frac{x^3}{3} \ln(x) - \frac{x^3}{9}$
4. $\int \sin(\ln(x)) dx = \frac{x}{2} (\sin(\ln(x)) - \cos(\ln(x)))$
5. $\int x \ln\left(\frac{1-x}{1+x}\right) dx = \frac{x^2-1}{2} \ln\left(\frac{1-x}{1+x}\right) - x$
6. $\int \frac{\ln^2(x)}{x^2} dx = -\frac{\ln^2(x) + 2\ln(x) + 2}{x}$
7. $\int \frac{\ln(\ln(x))}{x} dx = \ln(x) (\ln(\ln(x)) - 1)$
8. $\int x^2 \arctan(3x) dx = \frac{x^3}{3} \arctan(3x) - \frac{x^2}{18} + \frac{1}{162} \ln(9x^2 + 1).$

Exercice 4 *Intégration des fractions rationnelles*

1. $\int \frac{x+1}{(x^2+4x+5)^2} dx = -\frac{x+3}{2(x^2+4x+5)} - \frac{1}{2} \arctan(x+2)$
2. $\int \frac{dx}{x^3-2x^2+x} = \ln|x| - \ln|x-1| - \frac{1}{x-1}$
3. $\int \frac{dx}{(x^2-4x+3)(x^2+4x+5)} = \frac{1}{52} \ln|x-3| - \frac{1}{20} \ln|x-1| + \frac{1}{65} \ln(x^2+4x+5) + \frac{7}{130} \arctan(x+2)$
4. $\int \frac{x^3-1}{4x^3-x} dx = \frac{x}{4} + \frac{1}{16} \ln \left| \frac{x^{16}}{(2x-7)^7(2x+1)^9} \right|$
5. $\int \frac{2x-3}{(x^2-3x+2)^3} dx = -\frac{1}{2(x^2-3x+2)^2}$
6. $\int \frac{x^7+x^3}{x^{12}-2x^4+1} dx = \frac{1}{2} \ln|x^4-1| - \frac{1}{4} \ln|x^8+x^4-1| - \frac{1}{2\sqrt{5}} \ln \left| \frac{2x^4+1-\sqrt{5}}{2x^4+1+\sqrt{5}} \right|$
7. $\int \frac{x^2 dx}{(x-1)^{10}} = -\frac{1}{9(x-1)^9} - \frac{1}{4(x-1)^8} - \frac{1}{7(x-1)^7}$
8. $\int \frac{dx}{x^8+x^6} = \frac{1}{5x^5} + \frac{1}{3x^3} - \frac{1}{x} - \arctan(x)$

Exercice 5 *Intégration des fonctions trigonométriques*

1. $\int \cos^3(x) dx = \sin(x) - \frac{\sin^3(x)}{3}$
2. $\int \frac{\cos^5(x)}{\sin^3(x)} dx = \frac{\sin^2(x)}{2} - \frac{1}{2\sin^2(x)} - \ln(\sin^2(x))$
3. $\int \frac{dx}{\sqrt{\tan(x)}} = \frac{1}{2\sqrt{2}} \ln \left(\frac{\tan^2(x) - \sqrt{2}\tan(x) + 1}{\tan^2(x) + \sqrt{2}\tan(x) + 1} \right) - \frac{1}{\sqrt{2}} \arctan \left(\frac{\sqrt{2}\tan(x)}{\tan^2(x) - 1} \right)$
4. $\int \frac{dx}{3 + 5\cos(x)} = \frac{1}{4} \ln \left| \frac{\tan\left(\frac{x}{2}\right) - 2}{\tan\left(\frac{x}{2}\right) + 2} \right|$
5. $\int \frac{1 + \tan(x)}{1 - \tan(x)} dx = -\ln |\sin(x) + \cos(x)|$
6. $\int \frac{dx}{\cos(x) + 2\sin(x) + 3} = \arctan \left(1 + \tan\left(\frac{x}{2}\right) \right)$
7. $\int \frac{1 - \sin(x) + \cos(x)}{1 + \sin(x) - \cos(x)} dx = -x + 2 \ln \left| \frac{\tan\left(\frac{x}{2}\right)}{1 + \tan\left(\frac{x}{2}\right)} \right|$
8. $\int \frac{dx}{\sinh(x) \cosh^2(x)} = \ln \left| \tanh\left(\frac{x}{2}\right) \right| + \frac{1}{\cosh(x)}$
9. Calculer pour $n = 3$ et $n = 4$ les intégrales $I_n = \int \frac{dx}{\cos^n(x)}$. On a $I_4 = \tan(x) + \frac{\tan^3(x)}{3}$ et $I_3 = \frac{\sin(x)}{2\cos(x)} + \frac{1}{2} \ln \left| \tan(x) + \frac{1}{\sin(x)} \right|$.
10. $\int \tan^2\left(\frac{x}{2} + \frac{\pi}{4}\right) dx = 2 \tan\left(\frac{x}{2} + \frac{\pi}{4}\right) - x$.
11. $\int x \arctan(2x + 3) dx = \frac{(x^2 - 2) \arctan(2x + 3)}{2} + \frac{3 \ln(2x^2 + 6x + 5)}{8} - \frac{x}{4}$.